SINUS HEADACHE DAY NIGHT- acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride AmeriSource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

GNP - 1149 - 2019-1002

SINUS + HEADACHE DAY

Drug Facts

Active ingredients (in each day caplet)	Purpose
	Pain
Acetaminophen 325 mg	reliever/fever
	reducer
Dhorylonhuino IICl F ma	Nasal
Phenylephrine HCl 5 mg	decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - headache
 - sinus congestion and pressure
 - nasal congestion
 - minor aches and pains
- helps decongest sinus openings and passages
- promotes sinus drainage
- helps clear nasal passages
- temporarily reduces fever

SINUS + HEADACHE NIGHT

Drug Facts

Active ingredients (in each night caplet)	Purpose
	Pain
Acetaminophen 325 mg	reliever/fever
	reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dhamilanhuina IICl F ma	Nasal
Phenylephrine HCl 5 mg	decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - headache
 - sinus congestion and pressure

- nasal congestion
- runny nose and sneezing
- minor aches and pains
- helps decongest sinus openings and passages
- promotes sinus drainage
- helps clear nasal passages
- temporarily reduces fever

Warnings

Liver warning

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage In addition, when using Sinus + Headache Night:
- excitability may occur, especially in children

- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take Day and Night caplets at the same time
- do not take more than a total of 10 caplets in 24 hours

adults and	•	take 2 caplets every 4 hours
children	-	swallow whole; do not crush, chew,
12 years		or dissolve
and over		
children		1 1 .
under 12	•	ask a doctor
years		

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

Sinus + Headache Day

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C blue #1, flavor, hypromellose, lactose anhydrous, magnesium stearate, povidone, pregelatinized starch, propylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Sinus + Headache Night

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, flavor, hypromellose, iron oxide ochre, lactose anhydrous, magnesium stearate, povidone, pregelatinized starch, propylene glycol,

PRINCIPAL DISPLAY PANEL

GOOD NEIGHBOR PHARMACY

NDC 46122-413-60

DAY & NIGHT

Sinus + Headache

pain reliever/fever reducer, nasal decongestant, antihistamine*

For Adults

acetaminophen, phenylephrine hcl, chlorpheniramine maleate*

*Antihistamine in Nighttime only

Daytime (Non-Drowsy)

Sinus Headache, Sinus Pressure, Nasal Congestion

COOL TASTE


12 CAPLETS

Nighttime

Sinus Headache, Sinus Pressure, Nasal Congestion, Runny Nose*

COOL TASTE

8 CAPLETS







Sinus + Headache pain reliever/fever reducer sinus decongestant, antitistamine

For Adults - Day & Night



NDC 46122-413-60



Sinus + Headache

reliever/fever reducer, nasal decongestant, antihistamine

For Adults

acetaminophen, phenylephrine hol, chlorpheniramine maleate*

*Antihistamine in Nighttime only

Daytime (Non-Drowsy)

- Sinus Headache
- Sinus Pressure
- Nasal Congestion

Nighttime

- Sinus Headache
- Sinus Pressure
- Nasal Congestion
- Runny Nose*

COOL TASTE 8 CAPLETS

COOL TASTE 12 CAPLETS

starch, propylene glycol, stearic acid, titanium dioxide flavor, hypromellose, iron oxide ochre, lactose anhydrous, magnesium stearate, povidone, pregelatinized Sinus + Headache Night sosculiame potassium, colloidal silicon dioxide, croscarmellose sodium,

acid, titanium dioxide

magnesium stearate, povidone, pregelatinized starch, propytene glycol, sodium starch glycolate, stearic dioxide, croscarmellose sodium, D&C yellow #10, FD&C blue #1, flavor, hypromellose, lactose anhydrous, Inactive ingredients Sinus + Headache Day acesuliame potassium, colloidal sillcon

nund racts (continued)

8 ARE TORN OR BROKEN

 retain carton for complete product information and warmings ■ store between 20-25°C (68-77°F) in a dry place Uther information

> ■ 92K 9 doctor children under 12 years 12 years and over ■ swsllow whole – do not crush, chew, or dissolve ■ take 2 caplets every 4 hours adults and children

> > ■ qo not take more than a total of 10 caplets in 24 hours

do not take Day and Night caplets at the same time

DIRECTIONS - do not take more than directed (see overdose warming)

notice any signs or symptoms.

(1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not



Distributed By AmerisourceBergen 1300 Morris Drive Chesterbrook, PA 19087 Questions or Concerns? www.mygnp.com

Overdose warming: in case of overdose, det medical belo or contact a Poison Control Center right away keep out of reach of children.

It pregnant or breast-feeding, ask a health professional before use.

Luese conig pe sidus of a serious condition.

 uew symptoms occur ■ fever gets worse or lasts more than 3 days redness or swelling is present

■ bein or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if

 pe caretul when drwing a motor vehicle or operating machinery Svoid alcoholic drinks

 slcohol, sedatives, and tranquilizers may increase drowsiness quowsiness may occur

 excuspility may occur, especially in children in addition, when using Sinus + Headache Mght:

 do not exceed recommended dosage When using this product

 taking sedatives or tranquifizers taking the blood thinning drug warrarin Ask a doctor or pharmacist before use if you are

s prestring problem such as emphysema or chronic brouchtis

trouble unnating due to an enlarged prostate gland

■ diabetes thyroid disease

 pidy plood bressure ■ heart disease ■ liver disease

Ask a doctor before use if you have

■ if you have ever had an allergic reaction to this product or any of its ingredients taking this product.

drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

Orug Facts (continued)



SINUS HEADACHE DAY NIGHT

acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source)

NDC:46122-413

Packaging

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:46122-413-60	1 in 1 CARTON	03/01/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8

Part 1 of 2

ACETAMINOPHEN, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen and phenylephrine hydrochloride tablet, coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name Basis of Strength			
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg	
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)		
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)		
CROSCARMELLOSE SODIUM (UNII: M28 O L1HH48)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
ANHYDRO US LACTO SE (UNII: 3S Y5LH9 PMK)		
MAGNESIUM STEARATE (UNII: 70097M6130)		
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)		
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)		

Product Characteristics				
Color	green	Score	no score	
Shape	OVAL (capsule-shaped)	Size	17mm	
Flavor	MINT	Imprint Code	AAA;1114	
Contains				

ı	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Marketing Start Date	Marketing End Date	
OTC monograph final	part341		

Part 2 of 2

ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet, coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg	
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	

Product Characte	Product Characteristics				
Color	white (Off-White)	Score	no score		
Shape	OVAL (capsule-shaped)	Size	17mm		
Flavor	MINT	Imprint Code	AAA;1117		
Contains					

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	8 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph final	part341					
3.5 1 .	.•					
Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph final	part341	03/01/2018				

Labeler - AmeriSource Bergen (007914906)

Revised: 10/2019 AmeriSource Bergen